


Service Description

Lipids are essential metabolites that have many key cellular functions and which can be analysed to gain insight into the metabolic state of cells. The number of lipid molecules in a cell, collectively called the lipidome, is estimated to be in the tens to hundreds of thousands.

According to the classification system proposed by the Lipid Metabolites and Pathways Strategy (LIPID MAPS) project, lipids are divided into eight classes: fatty acyls (FA), glycerolipids (GL), glycerophospholipids (GP), sphingolipids (SP), sterol lipids (ST), prenol lipids (PR), saccharolipids (SL), and polyketides (PK), covering a total of 43,413 lipid molecular species.



The common structures and main functions of eight lipid classes

Sun T, et al., Mass spectrometry-based lipidomics in food science and nutritional health: A comprehensive review, (2020)

Lipidomics is a new branch of metabolomics, which analyzes the compositions and content changes of lipids (major categories, subclasses and molecular types) in biological samples such as cells, tissues, organs or body fluids. Mass Spec based lipidomics (LC-MS), involves the comparison of the lipidome between control and test groups in order to screen differential lipids by statistical analysis, so as to identify differences between lipid metabolism and physiological/pathological changes.


BGI has extensive experience in the field of lipidomics with well-developed reliable workflows using market leading technologies and a bioinformatics infrastructure that is second to none.

Research Applications



- Disease biomarkers research
- Pathogenesis and prognosis study on diseases
- Drug target research
- Animal special behavior mechanism and food/medicinal value research
- Plant growth and development research
- Plant disease resistance and insect resistance research
- Microbial drug resistance mechanism

Technology Platforms



Waters CSH C18 column



Waters ACQUITY UPLC




Thermo Q Exactive/Q Exactive HF-X

Service Advantages

State-of-the-art LC-MS/MS systems	Large scale and high volume sample experience	High-precision identification results	Strict quality control system
<ul style="list-style-type: none"> • Thermo Q Exactive/HF • Resolution up to 24,000, ensuring high spectral quality and accurate results 	<ul style="list-style-type: none"> • Sample preparation at a capacity of up to 1000+ per day • Large scale project experience with 1000s of samples 	<ul style="list-style-type: none"> • LipidSearch database (1.7 million lipid ions) • 100% identification is achieved through the standards • Identification credibility rating 	<ul style="list-style-type: none"> • Strict protocols governing the whole workflow • Double quality control process of isotopic internal standard and QC samples


Lipidomics Workflow




Bioinformatics Analysis Workflow

Standard:

1.1 Data Processing




1.2 Data Quality Control




1.3 Statistical and Relative Quantitative Analysis of Lipid Classes

1.4 Statistical Analysis and Screening of Differential Lipids



1.5 Differential Lipid Analysis




Customized Solution:

16S/Metagenome + lipidome correlation analysis


Transcriptome + lipidome correlation analysis

Proteome + lipidome correlation analysis


Examples of Data QC Analysis - Stability and Repeatability



Examples of Statistical and Relative Quantitative Analysis of Lipid Classes




Statistical Chart of Lipid Sub Classes




Detected Changes in Lipid Sub Class


Examples of Statistical Analysis of Differential Lipids




Score Graph of PLS-DA




Volcano Plot



Cluster Analysis



ROC Curve



Lipid Correlation Analysis

General Sample Requirements

SAMPLE TYPE	RECOMMENDED SAMPLE AMOUNT	MINIMUM SAMPLE AMOUNT
Serum, plasma	≥ 250 µL	≥ 50 µL
Urine	≥ 500 µL	≥ 50 µL
Animal and clinical tissues	≥ 200 mg	≥ 25 mg
Feces and intestinal contents	≥ 200 mg	≥ 50 mg
Cell	≥ 1×10 ⁷	≥ 5×10 ⁶
Microorganism	≥ 1×10 ⁷ or ≥ 100 mg	≥ 5×10 ⁶ or ≥ 25 mg
Culture medium, fermentation medium	≥ 1 mL	≥ 100 µL
Plant tissue	≥ 1 g	≥ 100 mg
Milk	≥ 1 mL	≥ 1 mL
Other body fluids (amniotic fluid, saliva, hemolymph, cerebrospinal fluid, etc.)	≥ 250 µL	≥ 50 µL

Turn Around Time

Sample size: 1-50, 3-5 weeks

To Learn More

To learn how your research can benefit from BGI's extensive experience in Lipidomics, visit www.bgi.com, write to us via info@bgi.com or contact your local BGI office.

International Head Offices

BGI Americas

One Broadway, 3rd Floor
Cambridge, MA 02142,
USA
Tel: +1 617 500-2741

BGI Europe

Ole Maaløes Vej 3,
DK-2200 Copenhagen N,
Denmark
Tel: +45 7026 0806

BGI Asia

16 Dai Fu Street,
Tai Po Industrial Estate,
New Territories, Hong Kong
Tel: +852 36103510

BGI Australia

L6, CBCRC, 300 Herston
Road, Herston, Brisbane,
Queensland 4006, Australia
Tel: +61 (07) 3362 0475



BGI Genomics



BGLGenomics

Services For Research Use Only. Not For Use in Diagnostic Procedures (unless specifically noted).

Copyright ©2021 BGI. The BGI logo is a trademark of BGI. All rights reserved.

All brand and product names are trademarks or registered trademarks of their respective holders.

Information, descriptions and specifications in this publication are subject to change without notice.

DNBSEQ™ is a trademark of MGI Co. Ltd.

DNBSEQ™ technology-based services are executed at our service laboratories in China.

Published: February 2021

Version:1.0

BGI
We Sequence, You Discover